

Table of Symptoms

Signs and Symptoms	Pre Manifest	Early	Middle	Late
Motor	Mild <u>walking difficulty</u>	Mild <u>chorea</u> , <u>bradykinesia</u> , abnormal eye movements and problems with walking.	Chorea, <u>dystonia</u> , <u>rigidity</u> , poor <u>co-ordination</u> and difficulty holding objects. <u>Balance</u> and walking problems which may result in falls. <u>Dysphagia</u> and <u>dysarthria</u> .	Severe bradykinesia and rigidity, dysarthria, dysphagia, <u>incontinence</u> , <u>chest complications</u> (e.g. aspiration, <u>altered posture</u>) and <u>difficulty with breathing control</u> (e.g. reduced endurance).
Cognitive	Difficulty with <u>complex thinking</u> tasks.	Mild <u>problems with planning</u> , <u>sequencing</u> , <u>organising</u> , and <u>prioritising</u> tasks.	<u>Intellectual decline</u> and <u>memory loss</u> . <u>Lack of insight</u> and self-awareness, <u>difficulty doing two things at the same time</u> (e.g. walking and talking)	<u>Dementia</u>
Behavioural/ Psychiatric	<u>Depression</u> , <u>Aggression</u> , <u>Irritability</u>	Sadness, Depression, Irritability	<u>Apathy</u> , repetition of tasks or thoughts, <u>impulsivity</u> , antisocial and suicidal behaviour, <u>paranoia</u> , <u>delusions or hallucinations</u> .	<u>Delirium</u>